

Recombinant allergen Alt a 1 for *Alternaria alternata* (Alternaria plant rot fungus)

CATALOG NUMBER: RAL0025

LOT NUMBER: #

RECOMBINANT ALLERGEN: Alt a 1 is the major allergen of *Alternaria alternata* (De Vouge *et al.*, 1996).

DESCRIPTION: the *Alternaria alternata* Alt a 1 has been prepared as the recombinant mature protein fused to a his-tag.

PRESENTATION: liquid protein solution

SOURCE: *Pichia pastoris*

MOLECULAR WEIGHT: determined by SDS-PAGE, the protein band is between molecular markers of 25,000 and 18,400 Da, while relative molecular mass calculated from amino acid sequence is 19,036.95 Da.

BATCH COMPOSITION:

COMPONENTS	COMPOSITION
his-Alt a 1	recombinant allergen with a his-tag
Storage buffer	20 mM phosphate buffer pH 7, 0.15 M NaCl and 0.1% polyoxyethylene (10) tridecyl ether

QUALITY CONTROL:

1. PROTEIN CONCENTRATION DETERMINED ESPECTROPHOTOMETRICALLY

DO₂₈₀ = 1.596
A_{0.1%} (=1 g/l) = 1.295
CONCENTRATION*: 1.23 mg/ml

* The measurement of the protein concentration has been performed with the theoretical extinction coefficient of the recombinant protein obtained from Gill and vonHippel, 1989. It is recommended that the users carry out their absorbance determinations to avoid equipment variabilities regarding final concentration, mainly in reproducibility analysis.

2. PURITY CONTROL IN SDS-PAGE: 15%

kDa Mw Alt a 1





Figure 1. SDS-PAGE analysis (15%) of 10 µl of recombinant allergen Alt a 1. Purity is >95% as determined by gel electrophoresis.

3. PROTEIN FINGERPRINT BY MASS SPECTROMETRY

Top Score: 88 for Alt a 1

Protein score is $-10 \cdot \log(P)$, where P is the probability that the observed match is a random event. Protein scores greater than 13 are significant ($p < 0.05$).



Concise Protein Summary Report

Format As Concise Protein Summary [Help](#)

Significance threshold $p < 0.05$

Max. number of hits 10

Re-Search All Search Unmatched

1. [Alt_a_1](#) Mass: 28226 Score: 88 Expect: 1.7e-009 Matches: 10
Alt_a_1

The MS was performed with a by MALDI TOF/TOF model UltrafleXtreme (Bruker).

4. ABSENCE OF PRECIPITATION AFTER A FREEZING AND THAWING CYCLE: ok

LOT SPECIFICATIONS:

1. CONCENTRATION: 1.23 mg/ml

2. TOTAL QUANTITY PER ALIQUOT: 1 mg

3. TOTAL VOLUME PER ALIQUOT: 0.853 ml

4. STORAGE: Protein is shipped with dry ice. Upon arrival, it should be aliquoted to avoid repeated freezing and thawing cycles and stored at -20°C to -80°C . In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation.

5. TESTED APPLICATIONS: none.

6. POSSIBLE APPLICATIONS: WB, DB, Indirect ELISA, positive control in direct ELISA, CLIA, lateral-flow. Where this product has not been tested for use in a particular technique, this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates. **We strongly recommend utilising anti-CCD (cross-reactive-carbohydrate determinants) absorbent to eliminate the anti-CCD IgE antibodies present in normal human sera to prevent cross-reaction of this glycoprotein. By using this absorbent, reactions resulting from CCDs are eliminated.**

7. OBSERVATIONS: proteins should be maintained frozen at high concentrations. In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation. Prior making test dilutions and after defrosting the protein, is recommended to remove possible protein aggregates by centrifuging the stock solution, avoiding alterations in the immobilization of the biomolecule to the solid surface.

RELATED PRODUCTS:

None.

BIBLIOGRAPHY:

De Vouge, M. W., Thaker, A. J., Curran I. H., Zhang, L., Muradia, G., Rode, H. and Vijay, H. M. Isolation and expression of a cDNA clone encoding an *Alternaria alternata* Alt a 1 subunit. 1996, *Int. Arch. Allergy. Immunol.* 111:385-95.

Gill SC, von Hippel PH. Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem.* 1989 Nov 1;182(2):319-26.

Important Notes: During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 µl or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the containers cap.

Although recombinant antigens are expressed in non-pathogenic *P. pastoris* and bacterial integrity is destroyed during purification, the antigen preparation should be handled as potentially infectious.

FOR RESEARCH AND COMMERCIAL USE *IN VITRO*: not for human *in vivo* or therapeutic use.