

## Recombinant antigen LipL32 for *Leptospira interrogans*

**CATALOG NUMBER:** RAG0063

**LOT NUMBER:** #

**RECOMBINANT ANTIGEN:** *Leptospira interrogans* lipL32 (Haake *et al.*, 2000)

**DESCRIPTION:** the recombinant antigen LipL32 has been prepared as the mature antigen fused to a his-tag. It is produced from the ORF of the lipL32 gene which codifies the mayor outer membrane protein of the spirochete *Leptospira interrogans* without the leader peptide.

**PRESENTATION:** liquid protein solution

**SOURCE:** *Pichia pastoris*

**MOLECULAR WEIGHT:** determined by SDS-PAGE, the protein band is at the molecular marker of 35,000 Da, while relative molecular mass calculated from amino acid sequence and without glycosylation is 30469.52 Da.

**BATCH COMPOSITION:**

COMPONENTS	COMPOSITION
his-lipL32	recombinant antigen with a his-tag in its C-terminus
Storage buffer	20 mM phosphate buffer pH 8 and 0.1 M KCl


**QUALITY CONTROL:**

**1. PROTEIN CONCENTRATION DETERMINED ESPECTROPHOTOMETRICALLY**

DO<sub>280</sub> = 1.23  
 A<sub>0.1%</sub> (=1 g/l) = 1.128  
 CONCENTRATION\*: 1.09 mg/ml

\* The measurement of the protein concentration has been performed with the theoretical extinction coefficient of the recombinant protein obtained from Gill and von Hippel, 1989

**2. PURITY CONTROL IN SDS-PAGE: 15%**



**Figure 1.** SDS-PAGE analysis (15%) of 10 µl of recombinant lipL32. Purity is >95% as determined by gel electrophoresis.

**3. ABSENCE OF PRECIPITATION AFTER A FREEZING AND THAWING CYCLE: ok**

**LOT SPECIFICATIONS:**

- 1. CONCENTRATION:** 1.09 mg/ml
- 2. TOTAL QUANTITY PER ALIQUOT:** 1 mg
- 3. TOTAL VOLUME PER ALIQUOT:** 0.962 ml
- 4. STORAGE:** Protein is shipped with dry ice. Upon arrival, it should be aliquoted in order to avoid repeated freezing and thawing cycles and stored at -20°C to -80°C.
- 5. APPLICATIONS:** not tested. Where this product has not been tested for use in a particular technique, this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates.
- 6. OBSERVATIONS:** Proteins should be maintained frozen at high concentrations. The dilution to be performed for ELISA assays should be made with a small quantity of protein, the same day of the experiment. In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation. Prior making test dilutions and after defrosting the protein, is recommended to remove possible protein aggregates by centrifuging the stock solution, avoiding alterations in the immobilization of the biomolecule to the solid surface.

**RELATED PRODUCTS:**

LipL32 (*E. coli*), LipL21.

**BIBLIOGRAPHY:**

**Haake, D., G. Chao, R. L. Zuerner, J. K. Barnett, D. Barnett, M. Mazel, J. Matsunaga, P. N. Levett, and C. A. Bolin.** The leptospiral major outer membrane protein LipL32 is a lipoprotein expressed during mammalian infection. *2000, Infect Immun.*, 68(4):2276–2285.

**Gill SC, von Hippel PH.** Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem.* 1989 Nov 1;182(2):319-26.

**Important Notes:** During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 µl or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the containers cap.

Although recombinant antigens are expressed in non-pathogenic *P. pastoris* and yeast integrity is destroyed during purification, the antigen preparation should be handled as potentially infectious.

**FOR RESEARCH AND COMMERCIAL USE *IN VITRO*: not for human *in vivo* or therapeutic use.**

**Recombinant antigen LipL32 for *Leptospira interrogans***

Rekom Biotech S.L. – BIC-Granada, Avda. Innovación, 1 – 18016, Granada (Spain) Tel: +34 958 63 70 85  
E-mail: [info@rekombiotech.com](mailto:info@rekombiotech.com) – Web: [www.rekombiotech.com](http://www.rekombiotech.com) – An ISO 9001 Certified Company

Page 2 of 2

