

IgG polyclonal antibody against CCD

CATALOG NUMBER: PAB0004

LOT NUMBER: #

IMMUNOGEN: Sorbent for anti-cross-reactive carbohydrate determinants (CCD) antibodies (Rekom Biotech catalog reference SOR0001)

SPECIFICITY: Monospecific, reacts only with CCD epitopes. Non-reactive with normal human serum.

PREPARATION: Isolated from sera of rabbits immunized with highly pure (>98%) sorbent CCD. Anti-CCD specific antibody was purified by protein G affinity chromatography.

PURITY: >95% (SDS-PAGE)

PRESENTATION: liquid protein solution

SOURCE: rabbit

BATCH COMPOSITION:

COMPONENTS	COMPOSITION
Storage buffer	20 mM phosphate buffer pH 7, 0.15 M NaCl and 0.1% sodium azide

LOT SPECIFICATIONS:

- 1. CONCENTRATION:** 2.16 mg/ml
- 2. TOTAL QUANTITY PER ALIQUOT:** 1 mg
- 3. TOTAL VOLUME PER ALIQUOT:** 0.463 ml
- 4. STORAGE:** Antibody is shipped with dry ice. Upon arrival, it should be aliquoted to avoid repeated freezing and thawing cycles and stored at -20°C to -80°C. In order to defrost the antibody, maintain the aliquot at 25°C without shaking to avoid aggregation.
- 5. WESTERN BLOT:** suggested titer approx. 1:10,000



Figure 1. Western blot analysis of the polyclonal Ab to detect CCD in a clarified cell lysate. Anti-CCD polyclonal Ab was used in a 1:9,000 dilution over 3 µg of yeast clarified extract.

6. ELISA ANALYSIS:

Antibody titration assay. A titration assay of the antibody was performed with a fixed sorbent immunogen concentration.




Figure 2. Titration of the polyclonal Ab diluted from 1/2,800 in a twofold series against constant sorbent (0.25 µg/ml). Mean values of OD are used.

The suggested titer for Ab PAB0004 is between 1/102,400 (0.02 µg/ml) and 1/204,800 (0.01 µg/ml).

7. OTHER TECHNIQUES: not tested. Where this product has not been tested for use in a particular technique, this does not necessarily exclude its use in such procedures.

Important Notes: Where this product has not been tested for use in a technique, this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates.

During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 µl or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the containers cap.

FOR RESEARCH AND COMMERCIAL USE *IN VITRO*: not for human *in vivo* or therapeutic use

IgG polyclonal antibody against CCD

Rekom Biotech S.L. – BIC-Granada, Avda. Innovación, 1 – 18016, Granada (Spain) – Tel: +34 958 63 70 85
E-mail: info@rekombiotech.com – Web: www.rekombiotech.com – An ISO 9001 and 13485 Certified Company

Page 2 of 2