

Recombinant antigen CFP10 for Mycobacterium tuberculosis

CATALOG NUMBER: RAG0050

LOT NUMBER: #

RECOMBINANT ANTIGEN: M. tuberculosis antigen CFP10 (Bekmurzayeva et al., 2013).

DESCRIPTION: the *Mycobacterium tuberculosis* gene for the 10 kDa culture filtrate antigen EsxB, has been expressed as a recombinant antigen fused to a his-tag in its N-terminus.

PRESENTATION: liquid protein solution

SOURCE: Escherichia coli

SPECIFIC **ANTIBODY** (CALIBRATOR): Polyclonal antibody for Tuberculosis (Rekom Biotech catalog reference PAB0009)

MOLECULAR WEIGHT: determined by SDS-PAGE, the protein band is between molecular markers of 25,000 and 18,400 Da, while relative molecular mass calculated from amino acid sequence is 17,506.0 Da.

BATCH COMPOSITION:

COMPONENTS	COMPOSITION
his-CFP10	recombinant antigen with a his-tag in its N-terminus
Storage buffer	20 mM phosphate buffer pH 6.5, 0.15 M NaCl and 0.1% polyoxyethylene (10) tridecyl ether

QUALITY CONTROL:

PROTEIN CONCENTRATION **DETERMINED ESPECTROPHOTOMETRICALLY**

 $DO_{280} = 1.65$

 $A_{0.1\%}$ (=1 g/l) = 0.484

CONCENTRATION*: 3.43 mg/ml

2. PURITY CONTROL IN SDS-PAGE: 15%




Figure 1. SDS-PAGE analysis (15%) of 2 μl of recombinant CFP10. Purity is >98% as determined by gel electrophoresis.

3. ABSENCE OF PRECIPITATION AFTER A FREEZING AND THAWING CYCLE: ok

LOT SPECIFICATIONS:

1. CONCENTRATION: 3.43 mg/ml

2. TOTAL QUANTITY PER ALIQUOT: 1 mg

3. TOTAL VOLUME PER ALIQUOT: 0.306 ml

- 4. STORAGE: Protein is shipped with dry ice. Upon arrival, it should be aliquoted to avoid repeated freezing and thawing cycles and stored at -20°C to -80°C. In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation.
- 5. TESTED APPLICATIONS: none.
- 6. POSIBLE APPLICATIONS: WB, DB, Indirect ELISA, positive control in direct ELISA, CLIA, lateral-flow. Where this product has not been tested for use in a particular technique, this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates.
- **7. OBSERVATIONS:** proteins should be maintained frozen at high concentrations. The dilution to be performed for ELISA assays should be made with a small quantity of protein, the same day of the experiment. In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation. Prior making test dilutions and after defrosting the protein, is recommended to remove possible protein aggregates by centrifuging the stock solution, avoiding alterations in the immobilization of the biomolecule to the solid surface.

RELATED PRODUCTS:

CFP10:ESAT6.

BIBLIOGRAPHY:

Bekmurzayeva A., Sypabekova M. and D. Kanayeva. Tuberculosis diagnosis using immunodominant, secreted antigens of Mycobacterium tuberculosis. 2013. Tuberculosis 93: 381-388.

Gill SC, von Hippel PH. Calculation of protein extinction coefficients from amino acid sequence data. Anal Biochem. 1989 Nov 1;182(2):319-26.



Page 1 of 2

The measurement of the protein concentration has been performed with the theoretical extinction coefficient of the recombinant protein obtained from Gill and vonHippel, 1989. It is recommended that the users carry out their absorbance determinations to avoid equipment variabilities regarding final concentration, mainly in reproducibility analysis.

rekom

Important Notes: During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 μ l or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the containers cap.

Although recombinant antigens are expressed in non-pathogenic *E. coli* and bacterial integrity is destroyed during purification, the antigen preparation should be handled as potentially infectious.

FOR RESEARCH AND COMMERCIAL USE IN VITRO: not for human in vivo or therapeutic use.

