

# Recombinant biotinylated multi-epitope chimeric antigen for *Treponema pallidum* (ChimSyphilis2)

**CATALOG NUMBER:** RAG0064BIOT

**LOT NUMBER:** #

**RECOMBINANT ANTIGEN:** *Treponema pallidum* multi-epitope chimeric antigen.

**DESCRIPTION:** a recombinant multi-epitope chimeric antigen has been prepared by expressing Tpp15 and TmpA proteins of the spirochete *Treponema pallidum*.

**PRESENTATION:** liquid protein solution

**SOURCE:** *Escherichia coli*

**MOLECULAR WEIGHT:** determined by SDS-PAGE, the protein band is between the molecular markers of 116,000 Da and 66,200 Da, while relative molecular mass calculated from amino acid sequence is 68,372.94 Da.

**BATCH COMPOSITION:**

COMPONENTS	COMPOSITION
ChimSyphilis2-his	recombinant chimeric antigen with a his-tag
Storage buffer	20 mM phosphate buffer pH 8, 0.15 M NaCl and 0.1% polyoxyethylene (10) tridecyl ether


**QUALITY CONTROL:**

**1. PROTEIN CONCENTRATION DETERMINED ESPECTROPHOTOMETRICALLY**

DO<sub>280</sub> = 0.66  
 A<sub>0.1%</sub> (=1 g/l) = 0.491  
 CONCENTRATION\*: 1.33 mg/ml

\* The measurement of the protein concentration has been performed with the theoretical extinction coefficient of the recombinant protein obtained from Gill and vonHippel, 1989. It is recommended that the users carry out their absorbance determinations to avoid equipment variabilities regarding final concentration, mainly in reproducibility analysis.

**2. PURITY CONTROL IN SDS-PAGE: 12%**




**Figure 1.** SDS-PAGE analysis (12%) of 3 µl of recombinant ChimSyphilis2. Purity is > 95% as determined by gel electrophoresis.

**3. DISCRIMINATION OF PRE-VALIDATED SERA BY AN INDIRECT ELISA ASSAY**


The cut-off has been suggested about an "in-house" ELISA kit performed in Rekom Biotech.

Each end-user should carry out an analysis for their particular application.



**Figure 2.** The dot plot graph illustrates the distribution of positive and negative sera by an indirect IgG ELISA with a ChimSyphilis2 plate coating of 1 µg/ml. Pre-validated sera by Dx (IgG, Abbott-ELISA, Becton Dickinson-RPR, Spin React - TPHA) were used in this analysis. The chart shows the optical density at 450/620 nm for positive (blue) and negative (grey) IgG sera.

**4. WESTERN BLOT WITH STREPTAVIDIN TO DETECT BIOTINYLATION**



**Figure 3.** Western blot analysis to detect streptavidin /biotin reaction. The incubation was performed with HRP conjugated streptavidin (1:2500)

**5. ABSENCE OF PRECIPITATION AFTER A FREEZING AND THAWING CYCLE: ok**

**LOT SPECIFICATIONS:**

**1. CONCENTRATION:** 1.33 mg/ml

**2. TOTAL QUANTITY PER ALIQUOT:** 1 mg

**3. TOTAL VOLUME PER ALIQUOT:** 0.789 ml

**4. SUGGESTED TITER BY ELISA:** up to 1:1,330, which corresponds to 1 µg/ml of protein concentration in plates for IgG detection in an indirect ELISA assay.

**5. STORAGE:** Protein is shipped with dry ice. Upon arrival, it should be aliquoted to avoid repeated freezing and thawing cycles and stored at -20°C to -80°C. In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation.

**6. TESTED APPLICATIONS:** ELISA.

**7. POSSIBLE APPLICATIONS:** WB, DB, Capture ELISA, Double Antigen Sandwich (DAS), nanoparticles binding, plate orientation. Where this product has not been tested for use in a particular technique, this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a

**8. OBSERVATIONS:** proteins should be maintained frozen at high concentrations. The dilution to be performed for ELISA assays should be made with a small quantity of protein, the same day of the experiment. In order to defrost the protein, maintain the aliquot at 25°C without shaking to avoid aggregation. Prior making test dilutions and after defrosting the protein, is recommended to remove possible protein aggregates by centrifuging the stock solution, avoiding alterations in the immobilization of the biomolecule to the solid surface.

**RECOMMENDED MATCHED ANTIGEN PAIRS:**

CAPTURE: RAG0064  
DETECTION: RAG0064BIOT

**RELATED PRODUCTS:**

TmpA, Tpp15, Tpp15-Biot, Tpp17, Tpp17-Biot, Tpp47, Tpp47-Biot, ChimSyphilis1, ChimSyphilis1-Biot, ChimSyphilis2.

**BIBLIOGRAPHY:**

**Gill SC, von Hippel PH.** Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem.* 1989 Nov 1;182(2):319-26.

**Important Notes:** During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 µl or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the containers cap.

Although recombinant antigens are expressed in non-pathogenic *E. coli* and bacterial integrity is destroyed during purification, the antigen preparation should be handled as potentially infectious.

**FOR RESEARCH AND COMMERCIAL USE *IN VITRO*: not for human *in vivo* or therapeutic use.**